

Exam 4 Test Prep Worksheet

Alkenes			
Reaction	Reagents	Regioselectivity	Stereoselectivity
Hydrohalogenation			
Hydrohalogenation with peroxide			
Acid-Catalyzed Hydration			
Oxymercuration-Demercuration			
Hydroboration- oxidation			
Halogenation			
Halohydrination			
Catalytic Hydrogenation			
Syn-Dihydroxylation			
Oxidative Cleavage with hot potassium permanganate			
Ozonolysis			


Alkenes			


Alkynes			
Reaction	Reagents	Regioelectivity	Stereoselectivity
Halogenation			
Hydrohalogenation			
Hydrogenation			
Hydrogenation with Lindlar's			
Ozonolysis			
Oxidative Cleavage with hot potassium permanganate			
Formation of Acetylide			
Dissolving Metal Reduction			


Alkynes			





Given the reactions below:





1. Name the reaction
2. Predict the major product of the reaction
3. Draw the mechanism (if needed)














Predict the major product of the multi-step problems below:







Provide the forward synthesis reagents for the problems below:

3-methyl-2-butanol to 2-methyl-2-butanol



Provide a retrosynthesis for the scenarios below:







Multiple-choice questions


1. Addition of Br₂ to (E)-hex-3-ene produces
 - A) a meso dibromide
 - B) a mixture of enantiomeric dibromides which is optically active
 - C) a mixture of enantiomeric dibromides which is optically inactive
 - D) (Z)-3,4-dibromo-3-hexene
 - E) (E)-3,4-dibromo-3-hexene
2. Which of the following additions to alkenes occur(s) specifically in an anti fashion?
 - A) hydroboration-oxidation
 - B) addition of Br₂
 - C) addition of H₂
 - D) addition of H₂O in dilute acid
 - E) both A and B
3. HBr can be added to an alkene in the presence of peroxides (ROOR). What function does the peroxide serve in this reaction?
 - A) nucleophile
 - B) electrophile
 - C) radical chain initiator
 - D) acid catalyst
 - E) solvent

¹H NMR


In the compounds below, circle and label the unique hydrogen environments, their splitting pattern, and integration











Which of the indicated protons in each pair will show up farther downfield?





or





or





or




or



or




or




Given the molecular formula and ^1H NMR spectra, propose a structure

C_6H_{12}








C_6H_{10}




¹³C NMR


Predict how many signals you would see in the ¹³C NMR spectrum of each of these molecules:



1. Which of options indicates the correct order of carbon chemical shifts of the four carbons of the following compound? Draw what the ¹³C NMR would look like.




- A C_{Me} < C2 < C3 < C1
B C3 < C2 < C_{Me} < C1
C C1 < C3 < C2 < C_{Me}
D C1 < C_{Me} < C2 < C3



Given the following molecular formula, ^1H NMR, ^{13}C NMR, answer the following questions:



$\text{C}_6\text{H}_{10}\text{O}_2$



1. Is there symmetry in this compound?
2. In the ^{13}C NMR, what does the chemical shift of the signal at about 105 suggest?
3. In the ^1H NMR, what does the chemical shift of the signal at about 5.2 suggest?
4. In the ^1H NMR, what does the integration of the signal at about 2.3 suggest?
5. In the ^1H NMR, what does the splitting pattern of the signal at about 1.0 suggest?
6. What is the structure of this unknown?

Mass Spectrometry

Below is the mass spectrum of 2-butanone. Explain the large peak at 43 and the small peak at 73.



NIST Chemistry WebBook (<https://webbook.nist.gov/chemistry>)

1. A prominent ($M^+ - 18$) peak suggests that the compound might be a(n):

- A) Alkane
- B) Alcohol
- C) Ether
- D) Ketone
- E) Primary amine

2. Mass spectrometry detects:

- I. Radicals
 - II. Radicals and radical cations
 - III. Radical cations
 - IV. Cations and Anions
- A) I and II
 - B) III only
 - C) IV only
 - D) I, II, and III
 - E) II and IV

3. What is the criteria for using mass spectrometry?

- A) To find the composition of the sample
- B) To find the relative mass of atoms
- C) To find the concentration of elements in the sample
- D) To find the properties of the sample